

UNCLASSIFIED

AD NUMBER
ADB261087
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Aug 2000. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott St., Fort Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, dtd 21 Feb 2003

THIS PAGE IS UNCLASSIFIED

AD _____

Award Number: DAMD17-96-1-6123

TITLE: Design, Sythesis, and Testing of Breast Cancer
Angiogenesis Inhibitors

PRINCIPAL INVESTIGATOR: Edward Schwartz, Ph.D.

CONTRACTING ORGANIZATION: Montefiore Medical Center
Bronx, New York 10467

REPORT DATE: August 2000

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S.
Government agencies only (proprietary information, Aug 00).
Other requests for this document shall be referred to U.S.
Army Medical Research and Materiel Command, 504 Scott Street,
Fort Detrick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are
those of the author(s) and should not be construed as an official
Department of the Army position, policy or decision unless so
designated by other documentation.

20010102 141

NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.

LIMITED RIGHTS LEGEND

Award Number: DAMD17-96-1-6123
Organization: Montefiore Medical Center
Location of Limited Rights Data (Pages):

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.

N. M. G. Allen Austin

12/8/00

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)**2. REPORT DATE**
August 2000**3. REPORT TYPE AND DATES COVERED**
Annual (15 Jul 99 - 14 Jul 00)**4. TITLE AND SUBTITLE**

Design, Sythesis, and Testing of Breast Cancer Angiogenesis Inhibitors

5. FUNDING NUMBERS

DAMD17-96-1-6123

6. AUTHOR(S)

Edward Schwartz, Ph.D.

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)Montefiore Medical Center
Bronx, New York 10467**E-MAIL:**

eschwart@aecom.yu.edu

**8. PERFORMING ORGANIZATION
REPORT NUMBER****9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)**U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012**10. SPONSORING / MONITORING
AGENCY REPORT NUMBER****11. SUPPLEMENTARY NOTES****12a. DISTRIBUTION / AVAILABILITY STATEMENT**

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government agencies only (proprietary information, Aug 00). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.

12b. DISTRIBUTION CODE**13. ABSTRACT (Maximum 200 Words)**

Thymidine phosphorylase (TP) catalyzes the phosphorolysis of thymidine and other pyrimidine 2'-deoxyribonucleosides. In addition, TP has been shown to possess angiogenic activity in a number of *in vitro* and *in vivo* assays, and its angiogenic activity has been linked to its catalytic activity. Several 5- and 6-substituted uracil derivatives were synthesized and evaluated for their ability to inhibit TP activity. In the past year we have identified a 6-amino-substituted uracil analog, 6-(2-aminoethyl)amino-5-chlorouracil (AEAC) to be one of the most active compounds. AEAC was found to be a competitive inhibitor of TP with a K_i of 165 nM. Human recombinant TP induced human umbilical vein endothelial cell (HUVEC) migration in a modified Boyden chamber assay *in vitro*, and this action could be abrogated by the AEAC. This was specific for TP, as the inhibitor had no effect on the chemotactic actions of vascular endothelial growth factor (VEGF). HUVEC migration was also induced when TP-transfected human breast carcinoma cells were used in a co-culture assay in place of the purified angiogenic factors, and a TP inhibitor nearly completely blocked the tumor cell-mediated migration. These studies suggest that inhibitors of TP may be useful in breast cancers which are dependent upon TP-driven angiogenesis.

14. SUBJECT TERMS

Breast Cancer, angiogenesis, thymidine phosphorylase

15. NUMBER OF PAGES

14

16. PRICE CODE**17. SECURITY CLASSIFICATION
OF REPORT**

Unclassified

**18. SECURITY CLASSIFICATION
OF THIS PAGE**

Unclassified

**19. SECURITY CLASSIFICATION
OF ABSTRACT**

Unclassified

20. LIMITATION OF ABSTRACT

Unlimited

Table of Contents

Cover page	1
Limited Legend	2
Report Document Page	3
Table of contents	4
Introduction	5
Body of report	6-12
Conclusion	12
References	13-14



Introduction. There is a need for alternative approaches to treat metastatic breast cancer. One rapidly developing area of investigation is the study of tumor angiogenesis, the process by which a growing tumor mass recruits the new blood vessels required for its continued growth, and through which the tumor can spread to distant sites. Indeed the neovascularization process is thought to be one of the rate limiting steps for the growth of primary and metastatic tumors (1-2). Most studies have demonstrated the importance of angiogenesis in the progression of human breast malignancies and found the extent of vascularization to be correlated with tumor size and an indicator of node metastasis (3-8). Several polypeptides and growth factors that are produced by breast cancer epithelial and stromal cells have been identified as having endothelial cell mitogenicity and angiogenic activity (1,9). Our studies are focusing on the angiogenic factor PD-ECGF, based on evidence demonstrating its role in experimental and human cancer, and the finding that PD-ECGF is identical to human thymidine phosphorylase (TP), an enzyme that catalyzes the reversible conversion of thymidine to thymine and 2'-deoxyribose-1-phosphate (10-12). When transfected into NIH 3T3 cells, TP was found to increase the vascularization of tumors growing in nude mice after sc inoculation (13). Similarly, overexpressing TP in MCF7 breast carcinoma cells markedly increased tumor growth *in vivo*, although it had no effect on the growth of the cells *in vitro* (14). Western blot analysis of primary human breast tissue showed that TP expression was elevated in the tumors compared to the normal tissue, a finding which provides clinical relevance to the transfection experiments. Studies suggest that the angiogenic and endothelial cell chemotactic activities of PD-ECGF are dependent upon its enzymatic activity, and this has been confirmed with site-directed mutagenesis studies (14,15-17). Of the angiogenic factors identified to date, TP is the only one in which an enzymatic activity of the factor is required for angiogenic activity. These observations serve as the basis for our hypothesis that inhibition of the catalytic activity of TP will also block its angiogenic properties. By synthesizing inhibitors of TP, we will be able to test this hypothesis and provide a basis for the development of a novel class of antitumor agents.

Technical objective 1: Synthesis of inhibitors of TP.

Our original proposal suggested the synthesis of four structural classes of pyrimidine and pyrimidine nucleosides for evaluation as potential TP inhibitors. Recent developments in the area of TP inhibitors as potential antiangiogenic agents which had taken place in the last 2-3 years led us to incorporate a fifth class of analogs in last year's report.


In the previous year, we have concentrated our synthetic efforts on the preparation of derivatives of class I and V (see below) since these two series had demonstrated the highest activities found as yet and seemed to hold the best promise for the generation of new potent analogs. Efforts were also expended on the resynthesis, at larger scales, of compounds previously synthesized for more extensive biological evaluation.

Figure 1




The most active members of these two classes of analogs discovered to date have been compiled in table 1. (See below). Compounds 2 – 12 (all derivatives of class I) were prepared as shown in Fig. 2 (a) by substitution of the 6-chloro group of a variety of uracil derivatives by appropriate primary amines of commercial sources of readily available through procedures reported in the chemical literature. The 6-chloro substituted pyrimidine precursors were themselves obtained by adaptations (or modifications) of published procedures.

Figure 2



The precursor 5,6-dichlorouracil was obtained by the chlorination of commercially available 6-chlorouracil with sulfuryl chloride in acetic acid (Fig.2 (b)). The 5-methyl substituted derivative was obtained by substitution of the 6-OH group of 5-methylbarbituric with Cl using phosphorous oxychloride in 85% phosphoric acid (Fig.2 (c)).





Figure 3














Compounds 13-20 (all derivatives of class II) were prepared as shown in Fig. 3 (a) by substituting the Cl group of variously 5-substituted 6-chloromethyluracils with the appropriate amine or thiono- derivative. Precursor 5-chloro-6-chloromethyluracil was obtained by direct chlorination of the 5-position of 6-chloromethyluracil (commercially available) with sulfuryl chloride in acetic acid (Fig.3 (b)). The 5-methyl-6-chloromethyluracil precursor was obtained from readily available 5,6-dimethyluracil by oxygenation to the 6-aldehyde derivative, reduction to the 6-hydroxymethyl intermediate and chlorination with thionyl chloride and dimethylformamide (Fig.3 (c)).

Technical objective 2: Evaluation of target compounds as TP inhibitors.

Table 1: Inhibition of TP activity by 5-, 6-substituted uracil analogs.

			
Compound	R ₁	R ₂	IC ₅₀
1	Br	-NH ₂	17 μM
2	Cl	-NHCH ₂ CH ₂ NH ₂ HCl	0.25 μM
3	CH ₃	-NHCH ₂ CH ₂ NH ₂ HCl	1.7 μM
4	Cl	-NHCH ₂ CH ₂ OH	100 μM
5	CH ₃	-NHCH ₂ CH ₂ OH	> 1000 μM
6	Cl	-NHCH ₂ CH ₂ CH ₂ OH	650 μM
7	H		30 μM
8	H		114 μM
9	H		122 μM


10	H		160 μ M
11	H		220 μ M
12	H		275 μ M
13	H		>300 μ M
14	H		10 μ M
15	Cl		0.30 μ M
16	Cl		500 μ M
17	CH ₃		200 μ M
18	Cl		15 μ M

19	H		> 300 μ M
20	Cl		300 μ M

TP activity was measured in assays which contained 0.2 M KH_2PO_4 (pH 7.8), 0.2 mM $[5\text{-}^3\text{H}]$ thymidine (1 μCi), human TP (5 ng), and several different concentrations of the inhibitor being tested. Reactions were stopped after 30 min at 37 by the addition of ice-cold TCA containing activated charcoal. After centrifugation, an aliquot of the supernatant was counted in a liquid scintillation counter. IC_{50} values for each compound shown in Table 1 were calculated from data from at least 3 determinations.

We next further the mechanism of inhibition of compound 2, 6-aminoethylamino-5-chlorouracil (AEAC) on TP inhibition by varying both the concentration of inhibitor, and that of the substrate thymidine.

FIG. 4: Double reciprocal analysis of TP inhibition by AEAC.



Assays of TP activity were carried out as described in Table 1 except that the concentration of substrate thymidine was varied between 33 and 200 μM . AEAC was used at concentrations of 0 mM (○), 0.3 mM (■), and 0.9 mM (●). Data are means of two experiments.


Conclusion: Compound 2 (AEAC) is a competitive inhibitor of TP.

We next examined the effect of AEAC and a second potent TP inhibitor, 5-chloro-6(1-imidazolyl-methyl) uracil (CIMU; compound 15) on TP-mediated endothelial cell migration.

Human umbilical vein endothelial cells (HUVEC) were isolated from umbilical cords obtained less than 5 hours after delivery. After rinsing, veins were incubated with collagenase (100 U/ml) in Medium 199 with Earles salts for 10 min. Primary cultures were seeded into flasks precoated with 0.02% (w/v) gelatin in PBS. Culture medium consisted of M199 containing 20% newborn calf serum, 5% pooled human serum, 2 mM L-glutamine, 5 U/ml penicillin and 5 µg/ml streptomycin. Cells were incubated at 37° in a humidified incubator with 5% CO₂ with a medium change after 24 hours and every 2 days thereafter until confluent. Primary cultures of HUVEC were harvested by incubation with 0.05% trypsin/0.02% EDTA, and the cells collected and cell number determined using a Coulter counter.

Modified Boyden chamber migration assay. HUVEC (passages 1-4) were harvested using cell dissociation solution (Sigma) and pelleted by centrifugation. Cells (10⁵) were placed into transwell inserts (Becton Dickinson; 8 µm pore) precoated with fibronectin (10 µg/ml). Inserts containing cells were placed into a 24 well plate containing 0.7 ml M199 with 1% serum and incubated (37°) for 1 hour. A chemotactic stimulus, recombinant human TP (200 ng/ml) or human VEGF (10 ng/ml; R&D Systems) was added to the lower wells in the presence and absence of various concentrations of inhibitors, and the cells incubated for 5 hours at 37°. At the end of the incubation period, the surface of the upper membrane was swabbed with a cotton-tipped applicator to remove non-migrating endothelial cells. Wells and inserts were washed 3 times with PBS and stained for 30 min. in medium with 10 µM Cell Tracker Green (Molecular Probes). The cluster plate was washed 3X with PBS, cells fixed in 3.7% formaldehyde for 10 min. and mounted on microscope slides. For quantitation, five random fields were photographed (100x total magnification) and the number of HUVEC in each field counted.

FIG. 5: Effect of TP inhibitors on endothelial cell migration.



Human endothelial cells, isolated as described above, were used in a modified Boyden chamber assay. Cells (10⁵) were placed into fibronectin-coated transwell inserts, which were then placed into 24 well plates in which the lower wells contained 0.7 ml M199 with 1% serum. After 1 hour, recombinant human TP (200 ng/ml) or human VEGF (10

ng/ml) were added to the lower wells, as indicated (single hatched bars). Control wells (open bars) had no added angiogenic factor. Wells also contained the indicated concentrations (0 to 300 μ M) of either CIMU (double hatched bars) or AEAC (solid bars). Cells were incubated for 5 hours at 37°, at which time non-migrating endothelial cells were removed from the upper surface of the membrane. Cells were stained with Cell Tracker Green, washed with PBS, fixed in formaldehyde, and mounted on microscope slides. Migration was quantitated by counting five random photographed (100x total magnification) fields, and the number of HUVEC in each field counted.

Technical objective 3: *In vivo* assays. We have had difficulty in establishing *in vivo* assays for assessing the activity of our TP inhibitors. While we can observe angiogenesis using a Matrigel plug in mice, the assay is not reproducible and is highly subjective, making it unlikely to be useful for the quantitative assessment of the activity of the inhibitors. We have also not been able to establish tumors in nude mice from a MCF7 xenograft, despite attempting a number of different modifications to the protocol.

As an alternative to these approaches, we have adapted the Boyden chamber assay with HUVECs to incorporate human mammary tumor cells in a co-culture assay. These assays utilized tumor cells in place of the purified angiogenic factors; no other angiogenic stimulus was added to the incubation. Wild type MCF7 breast carcinoma cells and cells which had been stably transfected with a human TP cDNA (MCF7/TPneo) were used. 10⁵ cells were added to the lower wells and were allowed to attach for 24 hours. The medium was then replaced with M199 medium with and without 0.3 mM of the inhibitor CIMU (see above), combined with HUVEC-containing transwells, and migration over a 5 hour period was analyzed as described above.

TABLE 2: Inhibition of HUVEC migration by a TP inhibitor in a co-culture assay.

<u>Tumor Cells</u>	<u>CIMU</u>	<u>HUVEC cells per field</u>	<u>Inhibition</u>
None	-	11 \pm 1.7	
MCF7/TPneo	-	115 \pm 9.0	
MCF7/TPneo	+	28 \pm 2.6	84%

Data are means \pm SEM.

Conclusion: Breast cancer cell-mediated angiogenesis can be blocked by a TP inhibitor.

REFERENCES

1. Folkman J and Shing Y (1992) Angiogenesis. *J. Biol. Chem.* 267:10931-10934.
2. Hayes DF (1994) Angiogenesis and breast cancer. *Hematology-Oncology Clinics of North America* 8:51-71.

3. Weidner N, Semple, JP, Welch WR and Folkman J (1991) Tumor angiogenesis and metastasis: correlation in invasive breast carcinoma. *N. Engl. J. Med.* 324:1-8.
4. Horak ER, Leek R, Klenk N, LeJeune S, Smith K, Stuart N, Greenall M, Stepniewska K and Harris AL (1993) Angiogenesis, assessed by platelet/endothelial cell adhesion molecule antibodies, as indicator of node metastases and survival in breast cancer. *Lancet* 340:1120-1124.
5. Toi M, Kashitani J and Tominaga T (1993) Tumor angiogenesis is an independent prognostic indicator in primary breast carcinoma. *Int. J. Cancer* 55:371-374.
6. Gasparini G, Weidner N, Bevilacqua P. et. al. (1994) Tumor microvessel density, p53 expression, tumor size, and peritumoral lymphatic vessel invasion are relevant prognostic markers in node-negative breast carcinoma. *J. Clin. Oncol.* 12:454-466.
7. Fox SB, Leek RD, Smith K, Hollyer J, Greenal M, Harris AI (1994) tumor angiogenesis in node-negative breast carcinomas- relationship with epidermal growth factor receptor, estrogen receptor, and survival. *Breast Cancer Res. Treat.* 29:109-116.
8. Craft PS and Harris AL (1994) Clinical prognostic significance of tumor angiogenesis. *Annals of Oncology* 5:305-311.
9. Hlatky L, Tsionou C, Hahnfeldt P and Coleman CN (1994) Mammary fibroblasts may influence breast tumor angiogenesis via hypoxia-induced vascular endothelial growth factor up-regulation and protein expression. *Cancer Res.* 54:6083-6086.
10. Moghaddam A and Bicknell R (1992) Expression of PD-ECGF factor in *E. coli* and confirmation of its thymidine phosphorylase activity. *Biochemistry* 31:12141-12146.
11. Furukawa T, Yoshimura A, Sumizawa T, Haraguchi M, Akiyama S-I, Fukui K, Ishizawa M and Yamada Y (1992) Angiogenic factor. *Nature* 356:668.
12. Sumizawa T, Furukawa T, Haraguchi M, Yoshimura A, Takeyasu A, Ishizawa M, Yamada Y and Akiyama S-I (1993) Thymidine phosphorylase activity associated with platelet-derived endothelial cell growth factor. *J. Biochem.* 114: 9-14.
13. Ishikawa F, Miyazono K, Hellman U, Drexler H, Wernstedt C, Hagiwara K, Usuki K, Takaku F, Risau W and Heldin C-H (1989) Identification of angiogenic activity and the cloning and expression of platelet-derived endothelial cell growth factor. *Nature* 338: 557-562.
14. Moghaddam A, Zhang H-T, Fan T-P D, Hu D-E, Lees VC, Turley H, Fox SB, Gatter KC, Harris AL and Bicknell R (1995) Thymidine phosphorylase is angiogenic and promotes tumor growth. *Proc. Nat. Acad. Sci.* 92:998-1002.
15. Haraguchi M, Miyadera K, Uemura K, Sumizawa T, Furukawa T, Yamada K, Akiyama S-I, and Yamada Y (1992) Angiogenic activity of enzymes. *Nature* 368:198-199.

16. Finnis C, Dodsworth N, Pollitt CE, Carr G and Sleep D (1993) Thymidine phosphorylase activity of platelet-derived endothelial cell growth factor is responsible for endothelial cell mitogenicity. *Eur. J. Biochem.* 212:201-210.
17. Miyadera K, Sumizawa T, Haraguchi M, Yoshida H, Konstanty W, Yamada Y and Akiyama S (1995) Role of thymidine phosphorylase activity in the angiogenic effect of platelet-derived endothelial cell growth factor/thymidine phosphorylase. *Cancer Res.* 55:1687-1690.



DEPARTMENT OF THE ARMY
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

21 Feb 03

MEMORANDUM FOR Administrator, Defense Technical Information
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,
VA 22060-6218


SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical reports written for this Command. Request the limited distribution statement for the enclosed accession numbers be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Kristin Morrow at DSN 343-7327 or by e-mail at Kristin.Morrow@det.amedd.army.mil.

FOR THE COMMANDER:

Encl


PHYLLIS M. RINEHART
Deputy Chief of Staff for
Information Management

ADB263458	ADB282838
ADB282174	ADB233092
ADB270704	ADB263929
ADB282196	ADB282182
ADB264903	ADB257136
ADB268484	ADB282227
ADB282253	ADB282177
ADB282115	ADB263548
ADB263413	ADB246535
ADB269109	ADB282826
ADB282106	ADB282127
ADB262514	ADB271165
ADB282264	ADB282112
ADB256789	ADB255775
ADB251569	ADB265599
ADB258878	ADB282098
ADB282275	ADB232738
ADB270822	ADB243196
ADB282207	ADB257445
ADB257105	ADB267547
ADB281673	ADB277556
ADB254429	ADB239320
ADB282110	ADB253648
ADB262549	ADB282171
ADB268358	ADB233883
ADB257359	ADB257696
ADB265810	ADB232089
ADB282111	ADB240398
ADB273020	ADB261087
ADB282185	ADB249593
ADB266340	ADB264542
ADB262490	ADB282216
ADB266385	ADB261617
ADB282181	ADB269116
ADB262451	
ADB266306	
ADB260298	
ADB269253	
ADB282119	
ADB261755	
ADB257398	
ADB267683	
ADB282231	
ADB234475	
ADB247704	
ADB258112	
ADB267627	